

AUTHOR INDEX

- ACE Inhibition in Progressive Renal Disease Study Group, 12
- Ackerson LM, 117
- Adachi JD, 98
- ADMIRAL Investigators, 89
- Ahnen D, 17
- Albanese A, 109
- Albert H, 33
- Allredge BK, 86
- Almeida OP, 26
- Alpert JS, 88
- Altman DG, 64
- Angelli G, 11
- Antman EM, 6, 76
- Ard J, 48
- Armstrong PW, 73
- Assessment of Implementation Strategies (ASSIST) Trial Collaborative Group, 35
- Assessment of the Safety and Efficacy of a New Thrombolytic Regimen (ASSENT)-3 Investigators, 43
- Asthma Clinical Research Network of the National Heart, Lung, and Blood Institute, 14, 15
- Attia MW, 37
- Bacaltchuk J, 106-107
- Badgett R, 62
- Bagnall AM, 61
- Baisden CE, 47
- Baker R, 28
- Balden E, 62
- Baldessarini RJ, 63
- Barkhof F, 21
- Barragan P, 89
- Bates ER, 41
- Baty PJ, 108
- Bedford H, 71
- Belknap S, 53
- Ben-Shlomo Y, 109
- Berger W, 16
- Berns JS, 12
- Bhandari M, A-11-12 (No. 1)
- Bilous R, 101
- Bisgaard H, 102
- Black PN, 54
- Blanchard AR, 110-111
- Blecker D, 48
- Bliss DZ, 23
- Blue L, 87
- Boogaerts M, 58
- Borzak S, 42-43
- Bosch X, 41
- Boulé NG, 100
- Boushey HA, 14
- Bow EJ, 58
- Bower P, 108
- Bozzetti F, 93
- Braga M, 93
- Brass LM, 96
- Brenner BM, 84
- Bröchner-Mortensen J, 82
- Brocklebank D, 110-111
- Buchan DA, 13
- Burgess S, 63
- Campbell FA, 18-19
- Campbell M, 85
- Canadian Task Force on Preventive Health Care, 91
- Cannon CP, 4
- Capes SE, 114
- Capewell S, 31
- Cappelleri JC, 101
- CAPRICORN Investigators, 7
- Caraballo D, 77
- Carroll D, 18-19
- Carruthers SG, 35
- Cates C, 110
- CHD Risk Prediction Group, 36
- Chernoff R, 28
- Cheskin LJ, 49
- Clarke WR, 83
- Clopidogrel in Unstable Angina to Prevent Recurrent Events Trial Investigators, 45
- Clopton P, 68
- Cody J, 85
- Cohen LS, 26
- Colford J, 105
- Collaborative Group on Hormonal Factors in Breast Cancer, 115
- Collaborative Study Group, 83
- Comi G, 21
- Conlin PR, 48
- Cook DJ, 2
- Cooper ME, 84
- Covington C, 72
- Cranney A, 98
- D'Agostino Sr. RB, 36
- Daly C, 85
- Danner RL, 90
- DASH Collaborative Research Group, 48
- Datta M, 66
- Davidson RA, 13
- De Jonghe B, 1
- De Louvois J, 71
- de Zeeuw D, 84
- Decker JM, 3
- Deeks JJ, 103
- deFilippi CR, 8
- Del Mar CB, 37
- del Zoppo G, 50
- Delaney-Black V, 72
- Demicheli V, 103
- Demopoulos LA, 4
- Devereaux PJ, A-11-12 (No. 1), A-11-13 (No. 2)
- DeVer KA, 28
- Devillé WL, 113
- Dexter PR, 74
- Didier D, 30
- Diemer M, 116
- Donepezil MSAD Study Investigators Group, 59
- Donnan GA, 50
- Douketis JD, 9, 114
- Ducharme F, 99
- Duhaut P, 10
- Dupas JL, 92
- Dvorak J, 65
- Early Goal-Directed Therapy Collaborative Group, 90
- Early Treatment of Multiple Sclerosis Study Group, 21
- Eikelboom JW, 9
- Eloniemi-Sulkava U, 60
- Evans J, 57
- Everitt H, 22
- Fahy JV, 14
- Feit F, 45
- Feldman H, 59
- Ferrara A, 117
- Filippi M, 21
- Fish D, 56
- Fishbane S, 83-85
- Fletcher R, 91
- Flores RH, 97
- Frachi LM, 102
- French Catheter Study Group in Intensive Care, 1
- Frid PJ, 72
- Fries W, 52
- Furukawa TA, 20
- Gattinoni L, 55
- Gauthier S, 59
- Gavazzi C, 93
- Gelb AM, 86
- Gelfand RA, 101
- Gelijns AC, 88
- Gerber RA, 101
- Gerstein HC, 34, 114
- Ghali WA, A-11-12 (No. 1)
- Gianotti L, 93
- Gibson N, 68
- Giugliano RP, 6
- Godfrey C, 108
- Godskesen M, 33
- Golish JA, 110-111
- Golliot F, 1
- Goodman S, 4-5
- Görtsche PC, 20
- Gravenstein S, 24
- Griffin S, 100
- Grimm III R, 46
- Gross PA, 103
- Grosser S, 14-15
- Grundt S, 36
- Gula LJ, 29
- GUSTO IV-ACS Investigators, 44

- GUSTO V Investigators, 42
 Guthrie E, 27
 Guyatt GH, A-11-12 (No. 1), 2, A-11-13 (No. 2), A-17 (No. 2)
- Haddad E, 100
 Halket S, 71
 Handoll HH, 112
 Hanlon P, 46
 Hatcher S, 64
 Hatzigeorgiou C, 116
 Havstad S, 90
 Hawkins M, A-15 (No. 1)
 Hawton K, 64
 Hay P, 106-107
 Haynes RB, A-11-13 (No. 2)
 Head J, 109
 Hecker J, 59
 Hennen J, 63
 Hentinen M, 60
 Herrmann HC, 5
 Heywood P, 108
 Hicks G, 99
 Hirsch CH, 60
 Hochberg MC, 97
 Holbrook A, 74
 Hoogerwerf WA, 95
 HOPE Study Investigators, 34
 House A, 27
 Howarth N, 30
 Hróbjartsson A, 20
 Hu LT, 57
 Hughes CM, 1
 Hunninghake GW, 70
 Hunsicker LG, 83
 Hunt D, 114
 Hurley DL, 81
 Hurwitz B, 22
- Incorvaia B, 104
 Investigators of the "Duree Optimale du Traitement AntiVitamines K" (DOTAVK) Study, 10
 Irbesartan in Patients with Type 2 Diabetes and Microalbuminuria Study Group, 82
 Ireys HT, 28
 Isaacs SM, 86
 Itraconazole Neutropenia Study Group, 58
- Jackson JL, 62
 Jafar TH, 12
 Jaffrey JB, 69
 Jagasia DH, 7
 Jamal S, 98
 Janowsky EC, 115
 Jefferson TO, 103
 Jesse RL, 29
 Joffe H, 26
 Jung HJ, 23
- Kahrilas PJ, 95
 Kakuma R, 99
 Kamm MA, 92
 Kapoor WN, 77
 Kapur N, 27
 Karter AJ, 117
 Katzenschlager R, 109
 Kearon C, 10-11
 Kellum JA, 3
 Kenny GP, 100
 Kent DM, 36
 Kernan WN, 96
 Kessels AG, 69
 Kim Y, 28
 King S, 32
 Kinzbrunner BM, 18-19
 Kitching AD, 89
 Klein JD, 37
 Klempner MS, 57
 Knorr B, 102
 Koenig E, 52
 Kontos MC, 29
 Koretz RL, 93
 Korula J, 23
 Kourides IA, 101
 Kowdley KV, 94
 Krause J, 49
 Kreder HJ, 112
 Krishnaswamy P, 68
 Kronzon I, 45
- Lacasse Y, 70
 Lancaster T, 73
 Landa M, 12
 Lang E, 87
 Laville M, 3
 Lawson FM, 73
 Lazarus SC, 14
 Lee E, 17
 Lees AJ, 109
 Lehnert H, 82
 Lemanske RF, 15
 Lewis EJ, 83
 Lindsay GM, 46
 Liu J, 117
 Llevadot J, 6
 Lonergan E, 105
 Louis-Simonet M, 77
 Lubien E, 68
 Luce BR, 75
 Luckmann R, 59
 Luthra G, 92
 Luxenburg J, 105
- MacIntyre K, 31
 Mackway-Jones K, 27
 Madhok R, 112
 Mahmud N, 92
 Malmberg K, 114
 Mann JF, 34
- Mannion AF, 65
 Manns BJ, A-11-12 (No. 1)
 Mariani L, 93
 Marrugat J, 41
 Massel D, 6, 29
 Mauger EA, 15
 Mazer B, 16
 Mazur LJ, 102
 McAlister FA, 73
 McCann DV, 31
 McGreer A, 24
 McHugh F, 46
 McLeod RS, 91
 McMahon MM, 81
 McMurray JJ, 87
 McNagny SE, 96
 McQueen RH, 29
 Meade M, 55
 Mehra MR, 87
 Meier FA, 37
 Mellor Clark J, 108
 Merrer J, 1
 Meyerhoff J, 56-57
 Milgrom H, 16
 Millard PS, 67
 Miñana J, 94
 Moher M, 35
 Molinterno DJ, 5
 Montalescot G, 89
 Montori VM, A-11-12 (No. 1)
 Moore TJ, 48
 Morgan JE, 106-107
 Morrow DA, 76
 Moskowitz AJ, 88
 Mukherjee D, 53
 Müller F, 52
 Munari L, 104
 Müntener M, 65
 Murray J, 21, 104
- Nadelman RB, 56
 Natanson C, 90
 National Health Technology Assessment Inhaler Review Group, 110-111
 Navickis RJ, 2
 Nayak A, 16
 Nelemans PJ, 69
 Nguyen B, 90
 Nietsch HH, 94
 Ninet J, 10
 Nissen SE, 53
 Norword P, 24
 Norkola IL, 60
 Nowakowski J, 56
- O'Malley PG, 62
 Opstelten W, 113
 Ornato JP, 29
 Ortiz J, 94
 Overhage JM, 74

- Palmer CS, 75
 Panisko DM, A-14 (No. 1)
 Parkinson's Disease Research Group of the
 United Kingdom, 109
 Parsons L, 76
 Parving H-H, 82
 Pasricha PJ, 92, 95
 Pathak P, 114
 Perkins S, 74
 Perlman M, 71
 Perrier A, 30
 Pesenti A, 55
 Peters Jr. PH, 24
 Philbrick JT, 9
 Phillips EJ, 58
 Pinede L, 10
 Poole PJ, 54
 Prabhakar M, 6
 Prandoni P, 11
 PROGRESS Collaborative Group, 51
- Quinlan DJ, 9
- Rabbani GH, 67
 Rabbat CG, 82
 Rahman MM, 66
 Ram FS, 111
 Randomized Evaluation of Mechanical
 Assistance for the Treatment of
 Congestive Heart Failure (REMATCH)
 Study Group, 88
 Ray J, 76
 Rees PJ, 99
 RENAAL Study Investigators, 84
 Rennie D, A-17 (No. 2)
 Rice GP, 104
 Richter JE, 17, 95
 Rivers E, 90
 Rivetti D, 103
 Rogers MA, 13
 Rosanio S, 8
 Rose EA, 88
 Rowland N, 108
- Santamaria MG, 11
 Santoro JE, 62
 Sarasin FP, 77
 Savik K, 23
 Scheidtmann K, 52
 Schmid CH, 12
 Schoenbaum M, 116
- Scholten RJ, 113
 Schwartz B, 75
 Schwartz DA, 70
 Selby JV, 117
 Sheikh A, 22
 Sherbourne C, 116
 Shivkumar K, 7
 Shoor S, 113
 Sigal RJ, 100
 Skull SA, 25
 Small D, 13
 Smith BJ, 14-15
 Smith DA, 34
 Smith RC, 61
 Smith S, 117
 Soares CN, 26
 Sontag SJ, 95
 Sood B, 72
 Sorkness CA, 15
 Sowden AJ, 61
 Spechler SJ, 17
 Stanbrook MB, 30
 Stewart S, 31
 Stott DJ, 52
 Straus SE, A-11-13 (No. 3)
 Sullivan LM, 36
 Sverkey LP, 48
 Symmons DPM, 97
- TACTICS—Thrombolysis in
 Myocardial Infarction 18
 Investigators, 4
 Taimela S, 65
 TARGET Investigators, 5
 Tatum JL, 29
 Teka T, 67
 Teo KK, 73
 Thacker HL, 26
 The European Collaborative Study, 32
 Tick Bite Study Group, 56
 TIME Investigators, 47
 Tirschwell D, 51
 Tognoni G, 55
 Tomkins GE, 62
 Tondo L, 63
 Toochi M, 8
 Topol EJ, 5, 53
 Townsend E, 64
 Tracy JK, 97
 Tramer MR, 18-19
 Trefigloio R, 107
- Triplehorn C, 67
- Uhari M, 25
 Unttizer J, 116
- Van den Bergh G, 81
 Vasbinder GB, 69
 Vermund SH, 66
 Villanueva C, 94
 Viscoli CM, 96
 Vogel EW, 65
- Wahed MA, 66
 Wang EE, 25
 Wardlaw JM, 50
 Warfarin Optimal Duration Italian Trial
 Investigators, 11
 Weekers F, 81
 Weintraub WS, 4
 Weise WJ, 69
 Weiss R, 89
 Welch V, 98
 Wells GA, 100
 Westergaard J, 33
 White JE, 111
 Whiting P, 61
 Wiebe S, 86
 Wiedemann HP, 54
 Wilkes MM, 2
 Wilkinson D, 105
 Wilson K, A-17 (No. 3)
 Wilson P, 36
 Winston DJ, 58
 Wirth A, 49
 Wittenberg O, 89
 Wouters P, 81
 Wright F, 52
 Wright J, 110-111
 Wright L, 35
 Wyer PC, 8
- Yamaguchi T, 50
 Yi Q, 34
 Young G, 33
 Yudkin P, 35
- Zaman B, 67
 Zangwill KM, 75
 Zaoutis T, 37
 Zimmerman MB, 70

A

- Acenocoumarol
 - 3 months and 1 year of oral anticoagulant therapy were equivalent for idiopathic proximal deep venous thrombosis, 11
- Acetates
 - Montelukast moderately decreased asthma symptoms in children with persistent asthma, 102
- Adrenal cortex hormones
 - Omalizumab reduced inhaled corticosteroid use and exacerbations in childhood allergic asthma, 16
- Adrenergic beta-agonists
 - Carvedilol reduced mortality and morbidity caused by myocardial infarction in patients with left ventricular dysfunction, 7
 - Salmeterol was not as effective as triamcinolone for persistent asthma, 14
 - Stopping triamcinolone led to treatment failures in patients with persistent asthma who were receiving salmeterol and triamcinolone, 15
- Albumins
 - Review: Albumin administration is not associated with excess mortality in acutely ill patients, 2
- Albuminuria
 - Cardiovascular events were increased in high cardiovascular risk patients with albuminuria, 34
- Albuterol
 - Salmeterol was not as effective as triamcinolone for persistent asthma, 14
 - Stopping triamcinolone led to treatment failures in patients with persistent asthma who were receiving salmeterol and triamcinolone, 15
- Alcohol drinking
 - Low-level prenatal alcohol exposure was associated with adverse behavioral outcomes in children at 6 to 7 years of age, 72
- Algorithm
 - Helical computed tomographic results indicated small-to-moderate changes in the probability of pulmonary embolism, 30
- Alzheimer disease
 - Donepezil improved the clinical state and quality of life in moderate-to-severe Alzheimer disease, 59
- Aminosalicylic acids
 - Olsalazine was not better than placebo in maintaining remission in inactive Crohn disease, 92
- Amphotericin B
 - Itraconazole was as effective as amphotericin B for fever and neutropenia in cancer and led to fewer adverse events, 58
- Analgesics, non-narcotic
 - Review: Cannabinoids and codeine have similar effects on pain relief, but cannabinoids commonly cause adverse psychotropic effects, 18
- Anemia
 - Review: Recombinant human erythropoietin decreases the need for blood transfusions and may delay dialysis in chronic renal failure, 85
- Angioplasty, transluminal, percutaneous coronary
 - Abciximab was more effective than tirofiban in preventing ischemic events in patients having coronary stenting, 5
 - An early invasive strategy reduced the incidence of major cardiac events in patients with unstable coronary syndromes, 4
- Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly, 47
- Review: Platelet glycoprotein IIb/IIIa blockers for PCI or acute coronary syndromes reduce death and MI but increase bleeding, 41
- Angiotensin-converting enzyme inhibitors
 - Combined therapy with indapamide and perindopril but not perindopril alone reduced the risk for recurrent stroke, 51
 - Review: Angiotensin-converting enzyme inhibitors reduce the progression of nondiabetic renal disease, 12
- Antibiotics
 - Prolonged antibiotic treatment did not relieve chronic symptoms in Lyme disease, 57
 - Review: Antibiotics modestly increase remission rates in acute bacterial conjunctivitis, 22
 - Single-dose doxycycline prevented Lyme disease after an Ixodes scapularis tick bite, 56
- Anticoagulants
 - 3 months and 1 year of oral anticoagulant therapy were equivalent for idiopathic proximal deep venous thrombosis, 11
 - Long-term and short-term oral anticoagulation therapies were equivalent for venous thromboembolism, 10
 - Review: Extended-duration prophylaxis with heparin prevents deep venous thrombosis in hip or knee replacement, 9
- Anticonvulsants
 - Benzodiazepines were as safe as and more effective than placebo for out-of-hospital status epilepticus, 86
- Antidepressive agents
 - Review: Antidepressants improve headache in patients with chronic headache, 62
 - Review: Antidepressants increase remission and clinical improvement in bulimia nervosa, 106
 - Review: Psychological treatment is as effective as antidepressants for bulimia nervosa, but a combination is best, 107
- Antiemetics
 - Review: Cannabinoids control chemotherapy-induced nausea and vomiting but increase the risk for side effects, 19
- Antifungal agents
 - Itraconazole was as effective as amphotericin B for fever and neutropenia in cancer and led to fewer adverse events, 58
- Anti-HIV agents
 - Progression of disease in HIV-infected children slowed after the first year of life, 32
- Antihypertensive agents
 - Review: Angiotensin-converting enzyme inhibitors reduce the progression of nondiabetic renal disease, 12
- Anti-inflammatory agents, steroidal
 - Salmeterol was not as effective as triamcinolone for persistent asthma, 14
 - Stopping triamcinolone led to treatment failures in patients with persistent asthma who were receiving salmeterol and triamcinolone, 15
- Antiviral agents
 - Oseltamivir was safe and effective for prophylaxis of influenza in the frail elderly, 24
- Anxiety
 - Support by another mother and a child-life specialist decreased anxiety in mothers of children with chronic illnesses, 28

Appetite depressants

Continuous and intermittent sibutramine were equally effective at 44 weeks for reducing weight in obese persons, 49

Arthralgia

More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery, 33

Arthritis

Review: Adding newer disease-modifying drugs or biological agents to methotrexate improved rheumatoid arthritis symptoms, 97

Review: Studies on the cardiovascular effects of selective COX-2 inhibitors show mixed results, 53

Arthroplasty, replacement

Review: Extended-duration prophylaxis with heparin prevents deep venous thrombosis in hip or knee replacement, 9

Aspirin

Clopidogrel plus aspirin was effective but increased bleeding in acute coronary syndromes without ST-segment elevation, 45

Asthma

Montelukast moderately decreased asthma symptoms in children with persistent asthma, 102

Omalizumab reduced inhaled corticosteroid use and exacerbations in childhood allergic asthma, 16

Review: Adding antileukotrienes to inhaled corticosteroids reduces exacerbations in symptomatic chronic asthma, 99

Review: Pressurized metered-dose inhalers are as effective as other hand-held inhalers for corticosteroid use in asthma, 110

Review: Pressurized metered-dose inhalers are as effective as other hand-held inhalers for β_2 -agonist bronchodilator use in asthma, 111

Salmeterol was not as effective as triamcinolone for persistent asthma, 14

Stopping triamcinolone led to treatment failures in patients with persistent asthma who were receiving salmeterol and triamcinolone, 15

Atrial natriuretic factor

B-natriuretic peptide levels had high sensitivity and specificity for detecting abnormal ventricular function, 68

B

Beclomethasone

Omalizumab reduced inhaled corticosteroid use and exacerbations in childhood allergic asthma, 16

Biopsy

Clinical and radiologic evaluation had moderate sensitivity and specificity for detecting idiopathic pulmonary fibrosis, 70

Bipolar disorder

Review: Long-term lithium treatment lowers suicide risk in major affective disorder, 63

Blood glucose

Intensive insulin therapy reduced mortality and morbidity in critically ill patients, 81

Blood pressure monitoring

Home monitoring service was more effective than usual care in patients with essential hypertension, 13

Body mass index

Review: Exercise training reduces HbA_{1c} levels but not body mass in type 2 diabetes mellitus, 100

Brain ischemia

Review: Thrombolysis increases short-term death and intracranial hemorrhage but decreases long-term death or dependency, 50

Breast neoplasms

Review: Breast cancer is associated with a family history of the disease in first-degree relatives, 115

Bromocriptine

Initial bromocriptine did not change mortality in early, mild Parkinson disease, 109

Bronchitis

Review: Mucolytic drugs reduce exacerbations, illness days, and antibiotic use in chronic bronchitis and COPD, 54

Bronchodilator agents

Review: Pressurized metered-dose inhalers are as effective as other handheld inhalers for β_2 -agonist bronchodilator use in asthma, 111

Bulimia

Review: Antidepressants increase remission and clinical improvement in bulimia nervosa, 106

Review: Psychological treatment is as effective as antidepressants for bulimia nervosa, but a combination is best, 107

Bundle-branch block

Combining electrocardiography and necrosis biomarkers improved detection of acute myocardial infarction in patients with LBBB, 29

C

Cannabinoids

Review: Cannabinoids and codeine have similar effects on pain relief, but cannabinoids commonly cause adverse psychotropic effects, 18

Review: Cannabinoids control chemotherapy-induced nausea and vomiting but increase the risk for side effects, 19

Captopril

Review: CT angiography and magnetic resonance imaging are the best less-invasive tests for renal artery stenosis, 69

Cardiotonic agents

Review: Low-dose dopamine does not prevent acute renal failure or reduce mortality or need for hemodialysis, 3

Cardiovascular diseases

Cardiovascular events were increased in high cardiovascular risk patients with albuminuria, 34

Review: Studies on the cardiovascular effects of selective COX-2 inhibitors show mixed results, 53

Catheterization, peripheral

Femoral venous catheterization was associated with increased risk for infection and thrombosis in critically ill patients, 1

Ceftriaxone

Prolonged antibiotic treatment did not relieve chronic symptoms in Lyme disease, 57

Cerebrovascular accident

Combined therapy with indapamide and perindopril but not perindopril alone reduced the risk for recurrent stroke, 51

Estrogen did not prevent death or nonfatal stroke in postmenopausal women with ischemic stroke or TIA, 96

Levodopa plus carbidopa before physiotherapy increased motor recovery after stroke, 52

Review: Stress hyperglycemia after ischemic stroke indicates a greater risk for death in patients without diabetes, 114

- Review: Thrombolysis increases short-term death and intracranial hemorrhage but decreases long-term death or dependency, 50
- Chest pain
PredischARGE coronary angiography was better than exercise testing for reducing hospital use after low-risk chest pain, 8
- Chickenpox vaccine
Review: Varicella vaccination is effective in children, 25
- Child behavior disorders
Low-level prenatal alcohol exposure was associated with adverse behavioral outcomes in children at 6 to 7 years of age, 72
- Cholinesterase inhibitors
Donepezil improved the clinical state and quality of life in moderate-to-severe Alzheimer disease, 59
- Chronic disease
Support by another mother and a child-life specialist decreased anxiety in mothers of children with chronic illnesses, 28
- Cognition
Donepezil improved the clinical state and quality of life in moderate-to-severe Alzheimer disease, 59
- Colonoscopy
Review: Hemoccult screening reduces death from colorectal cancer in average-risk patients > 50 years of age, 91
- Colorectal neoplasms
Review: Hemoccult screening reduces death from colorectal cancer in average-risk patients > 50 years of age, 91
- Complementary therapies
The Desktop Guide to Complementary and Alternative Medicine: An Evidence-based Approach (Resource Corner), A-17 (No.3)
- Conjunctivitis, bacterial
Review: Antibiotics modestly increase remission rates in acute bacterial conjunctivitis, 22
- Continuity of patient care
Recall to a general practitioner or to a nurse clinic improved assessment in patients with coronary artery disease, 35
- Coronary angiography
PredischARGE coronary angiography was better than exercise testing for reducing hospital use after low-risk chest pain, 8
- Coronary artery bypass
A nurse-led intervention reduced risk factors, anxiety, and depression in patients waiting for CABG, 46
Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly, 47
- Coronary disease
Abciximab was more effective than tirofiban in preventing ischemic events in patients having coronary stenting, 5
The Framingham scores overestimated the risk for coronary heart disease in Japanese, Hispanic, and Native American cohorts, 36
PredischARGE coronary angiography was better than exercise testing for reducing hospital use after low-risk chest pain, 8
Recall to a general practitioner or to a nurse clinic improved assessment in patients with coronary artery disease, 35
Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly, 47
Review: Multidisciplinary CHD management programs improve the process of care and reduce hospitalizations, 73
- Counseling
Review: Primary care counseling improves psychological symptoms in patients with psychological and psychosocial problems, 108
- Creatine kinase
Combining electrocardiography and necrosis biomarkers improved detection of acute myocardial infarction in patients with LBBB, 29
- Critical care
Intensive insulin therapy reduced mortality and morbidity in critically ill patients, 81
Review: Albumin administration is not associated with excess mortality in acutely ill patients, 2
- Crohn disease
Olsalazine was not better than placebo in maintaining remission in inactive Crohn disease, 92
- Cyclooxygenase inhibitors
Review: Studies on the cardiovascular effects of selective COX-2 inhibitors show mixed results, 53
- ## D
- Decision support systems, clinical
Rule-based computerized reminders increased ordering of preventive services in an inpatient setting, 74
- Dementia
A community support program for dementia delayed but did not reduce institutionalization over 2 years, 60
Review: Haloperidol does not reduce agitation in dementia, 105
- Depression
1 of 2 quality-improvement interventions for depression in managed care was more effective but more costly than usual care, 116
17 β -estradiol reduced depressive and somatic symptoms in perimenopausal women, 26
- Developmental disabilities
Children infected with meningitis before 1 year of age were at increased risk for disability at 5 years, 71
- Diabetes mellitus
Cardiovascular events were increased in high cardiovascular risk patients with albuminuria, 34
Evidence-Based Diabetes Care (Resource Corner), A-15 (No. 1)
Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure, 83
Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria, 82
Losartan was renoprotective in diabetic nephropathy independent of its effect on blood pressure, 84
Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes, 101
Review: Exercise training reduces HbA_{1c} levels but not body mass in type 2 diabetes mellitus, 100
Review: Stress hyperglycemia after ischemic stroke indicates a greater risk for death in patients without diabetes, 114
Several simple rules predicted complications in high-risk patients with diabetes, 117
- Diarrhea, infantile
A rice-based diet with green banana or pectin reduced diarrhea in infants better than a rice diet alone, 67
Zinc or vitamin A reduced diarrhea in young, poor Bangladeshi children, 66
- Diazepam
Benzodiazepines were as safe as and more effective than placebo for out-of-hospital status epilepticus, 86

Diet

- The Dietary Approaches to Stop Hypertension diet lowered systolic blood pressure in stage-1 isolated systolic hypertension, 48
- A rice-based diet with green banana or pectin reduced diarrhea in infants better than a rice diet alone, 67

Dietary fiber

- Dietary fiber supplementation with psyllium or gum arabic reduced fecal incontinence in community-living adults, 23

Disease progression

- Progression of disease in HIV-infected children slowed after the first year of life, 32

Diuretics

- Combined therapy with indapamide and perindopril but not perindopril alone reduced the risk for recurrent stroke, 51

Dopamine agents

- Levodopa plus carbidopa before physiotherapy increased motor recovery after stroke, 52
- Review: Low-dose dopamine does not prevent acute renal failure or reduce mortality or need for hemodialysis, 3

Double-blind method

- "Double blind, you are the weakest link—good-bye!" (Editorial), A-11 (No. 1)

Doxycycline

- Prolonged antibiotic treatment did not relieve chronic symptoms in Lyme disease, 57
- Single-dose doxycycline prevented Lyme disease after an *Ixodes scapularis* tick bite, 56

E

Emergency medical services

- Benzodiazepines were as safe as and more effective than placebo for out-of-hospital status epilepticus, 86

Endoscopy

- Nadolol and isosorbide prevented recurrent variceal bleeding better than did endoscopic ligation in cirrhosis, 94

Enteral nutrition

- Enteral nutrition led to fewer postoperative complications than did parenteral feeding in gastrointestinal cancer, 93

Erythropoietin, recombinant

- Review: Recombinant human erythropoietin decreases the need for blood transfusions and may delay dialysis in chronic renal failure, 85

Esophageal and gastric varices

- Nadolol and isosorbide prevented recurrent variceal bleeding better than did endoscopic ligation in cirrhosis, 94

Esophageal neoplasms

- Surgery relieved symptoms but decreased survival more than medical treatment in gastroesophageal reflux disease, 17

Esophageal stenosis

- Surgery relieved symptoms but decreased survival more than medical treatment in gastroesophageal reflux disease, 17

Esophagitis

- Lansoprazole provided more effective and faster relief for heartburn than did omeprazole in erosive esophagitis, 95
- Surgery relieved symptoms but decreased survival more than medical treatment in gastroesophageal reflux disease, 17

Estradiol

- 17 β -estradiol reduced depressive and somatic symptoms in perimenopausal women, 26

- Estradiol did not prevent death or nonfatal stroke in postmenopausal women with ischemic stroke or TIA, 96

Estrogen replacement therapy

- Estrogen did not prevent death or nonfatal stroke in postmenopausal women with ischemic stroke or TIA, 96

Etidronic acid

- Review: Etidronate decreases vertebral but not nonvertebral fractures in postmenopausal women, 98

Evidence-based medicine

- Clinical expertise in the era of evidence-based medicine and patient choice (Editorial), A-11 (No. 2)
- The Desktop Guide to Complementary and Alternative Medicine: An Evidence-based Approach* (Resource Corner), A-17 (No.3)
- "Double blind, you are the weakest link—good-bye!" (Editorial), A-11 (No. 1)
- Evidence-Based Diabetes Care* (Resource Corner), A-15 (No. 1)
- Evidence-Based Hypertension* (Resource Corner), A-14 (No. 1)
- Evidence-based medicine in practice (Practice Corner), A-11 (No. 3)
- Users' Guides to the Medical Literature: A Manual for Evidence-Based Clinical Practice* (Resource Corner), A-14 (No. 2)

Exercise test

- Predischarge coronary angiography was better than exercise testing for reducing hospital use after low-risk chest pain, 8

Exercise

- Physiotherapy, aerobics, and training devices reduced pain intensity and frequency in chronic low-back pain, 65
- Review: Exercise training reduces HbA_{1c} levels but not body mass in type 2 diabetes mellitus, 100

Expectorants

- Review: Mucolytic drugs reduce exacerbations, illness days, and antibiotic use in chronic bronchitis and COPD, 54

F

Family practice

- 1 of 2 quality-improvement interventions for depression in managed care was more effective but more costly than usual care, 116
- Recall to a general practitioner or to a nurse clinic improved assessment in patients with coronary artery disease, 35

Family

- Support by another mother and a child-life specialist decreased anxiety in mothers of children with chronic illnesses, 28

Fatigue syndrome, chronic

- Review: Behavioral interventions show the most promise for the chronic fatigue syndrome, 61

Fecal incontinence

- Dietary fiber supplementation with psyllium or gum arabic reduced fecal incontinence in community-living adults, 23

Femoral vein

- Femoral venous catheterization was associated with increased risk for infection and thrombosis in critically ill patients, 1

Fibrinolytic agents

- Reteplase plus abciximab and reteplase alone led to similar 30-day mortality rates in acute MI, 42
- Review: Some new bolus tPA-derived fibrinolytics are as effective as tPA in acute MI, but others increase bleeding, 6
- Tenecteplase plus enoxaparin or abciximab was better than tenecteplase plus unfractionated heparin for acute MI, 43

Fluid therapy

- Review: Albumin administration is not associated with excess mortality in acutely ill patients, 2

Fractures

Review: Etidronate decreases vertebral but not nonvertebral fractures in postmenopausal women, 98

Frail elderly

Oseltamivir was safe and effective for prophylaxis of influenza in the frail elderly, 24

G

Gastroesophageal reflux

Surgery relieved symptoms but decreased survival more than medical treatment in gastroesophageal reflux disease, 17

Gastrointestinal neoplasms

Enteral nutrition led to fewer postoperative complications than did parenteral feeding in gastrointestinal cancer, 93

Glucocorticoids, synthetic

Review: Pressurized metered-dose inhalers are as effective as other hand-held inhalers for corticosteroid use in asthma, 110

H

Haloperidol

Review: Haloperidol does not reduce agitation in dementia, 105

Headache

Review: Antidepressants improve headache in patients with chronic headache, 62

Heart failure, congestive

Left ventricular assist devices reduced the risk for death and increased 1-year survival in chronic end-stage heart failure, 88

A specialist-nurse intervention reduced readmissions in patients hospitalized with chronic heart failure, 87

Heart-assist devices

Left ventricular assist devices reduced the risk for death and increased 1-year survival in chronic end-stage heart failure, 88

Heartburn

Lansoprazole provided more effective and faster relief for heartburn than did omeprazole in erosive esophagitis, 95

Hemoglobin A_{1c}, glycosylated

Review: Exercise training reduces HbA_{1c} levels but not body mass in type 2 diabetes mellitus, 100

HIV infections

Progression of disease in HIV-infected children slowed after the first year of life, 32

Home care services

A community support program for dementia delayed but did not reduce institutionalization over 2 years, 60

Hospital mortality

Mortality did not vary with sex by 1 year after a first acute myocardial infarction, 31

Hyperglycemia

Review: Stress hyperglycemia after ischemic stroke indicates a greater risk for death in patients without diabetes, 114

Hypertension

The Dietary Approaches to Stop Hypertension diet lowered systolic blood pressure in stage-1 isolated systolic hypertension, 48

Evidence-Based Hypertension (Resource Corner), A-14 (No. 1)

Home monitoring service was more effective than usual care in patients with essential hypertension, 13

Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure, 83

Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria, 82

Losartan was renoprotective in diabetic nephropathy independent of its effect on blood pressure, 84

I

IgE

Omalizumab reduced inhaled corticosteroid use and exacerbations in childhood allergic asthma, 16

Indapamide

Combined therapy with indapamide and perindopril but not perindopril alone reduced the risk for recurrent stroke, 51

Influenza vaccine

An intranasal influenza vaccine for the prevention of influenza in healthy children was cost-effective, 75

Oseltamivir was safe and effective for prophylaxis of influenza in the frail elderly, 24

Review: Inactivated vaccines provide the greatest protection against influenza in healthy persons, 103

Insulin

Intensive insulin therapy reduced mortality and morbidity in critically ill patients, 81

Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes, 101

Interferons

Interferon β -1a decreased the risk for conversion to clinically definite multiple sclerosis, 21

Review: In relapsing-remitting multiple sclerosis, recombinant interferon reduces exacerbations in the first 2 treatment years, 104

Isosorbide dinitrate

Nadolol and isosorbide prevented recurrent variceal bleeding better than did endoscopic ligation in cirrhosis, 94

Itraconazole

Itraconazole was as effective as amphotericin B for fever and neutropenia in cancer and led to fewer adverse events, 58

K

Kidney diseases

Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria, 82

Losartan was renoprotective in diabetic nephropathy independent of its effect on blood pressure, 84

Review: Low-dose dopamine does not prevent acute renal failure or reduce mortality or need for hemodialysis, 3

Review: Recombinant human erythropoietin decreases the need for blood transfusions and may delay dialysis in chronic renal failure, 85

L

Lansoprazole

Lansoprazole provided more effective and faster relief for heartburn than did omeprazole in erosive esophagitis, 95

Left ventricular function

Carvedilol reduced mortality and morbidity caused by myocardial infarction in patients with left ventricular dysfunction, 7

Leukotriene antagonists

Montelukast moderately decreased asthma symptoms in children with persistent asthma, 102

- Review: Adding antileukotrienes to inhaled corticosteroids reduces exacerbations in symptomatic chronic asthma, 99
- Levodopa
Initial bromocriptine did not change mortality in early, mild Parkinson disease, 109
Levodopa plus carbidopa before physiotherapy increased motor recovery after stroke, 52
- Lithium
Review: Long-term lithium treatment lowers suicide risk in major affective disorder, 63
- Lorazepam
Benzodiazepines were as safe as and more effective than placebo for out-of-hospital status epilepticus, 86
- Losartan
Irbesartan reduced progression of nephropathy caused by type 2 diabetes independent of the effect on blood pressure, 83
Losartan was renoprotective in diabetic nephropathy independent of its effect on blood pressure, 84
- Low back pain
Physiotherapy, aerobics, and training devices reduced pain intensity and frequency in chronic low-back pain, 65
- Lung diseases, obstructive
Review: Mucolytic drugs reduce exacerbations, illness days, and antibiotic use in chronic bronchitis and COPD, 54
- Lyme disease
Prolonged antibiotic treatment did not relieve chronic symptoms in Lyme disease, 57
Single-dose doxycycline prevented Lyme disease after an *Ixodes scapularis* tick bite, 56
- ### M
- Magnetic resonance angiography
Review: CT angiography and magnetic resonance imaging are the best less-invasive tests for renal artery stenosis, 69
- Maternal exposure
Low-level prenatal alcohol exposure was associated with adverse behavioral outcomes in children at 6 to 7 years of age, 72
- Medical records systems, computerized
Several simple rules predicted complications in high-risk patients with diabetes, 117
- Meningitis, bacterial
Children infected with meningitis before 1 year of age were at increased risk for disability at 5 years, 71
- Menisci, tibial
Review: Physical diagnostic tests have low diagnostic accuracy for meniscal lesions of the knee, 113
- Menopause
17 β -estradiol reduced depressive and somatic symptoms in perimenopausal women, 26
- Methotrexate
Review: Adding newer disease-modifying drugs or biological agents to methotrexate improved rheumatoid arthritis symptoms, 97
- Monitoring, physiologic
Early goal-directed therapy reduced mortality and multiorgan dysfunction in severe sepsis or septic shock, 90
- Motor skills
Levodopa plus carbidopa before physiotherapy increased motor recovery after stroke, 52
- Multiple sclerosis
Interferon β -1a decreased the risk for conversion to clinically definite multiple sclerosis, 21
Review: In relapsing-remitting multiple sclerosis, recombinant interferon reduces exacerbations in the first 2 treatment years, 104
- Myocardial infarction
Abciximab did not reduce death or myocardial infarction in the acute coronary syndrome without early revascularization, 44
Abciximab was more effective than tirofiban in preventing ischemic events in patients having coronary stenting, 5
Carvedilol reduced mortality and morbidity caused by myocardial infarction in patients with left ventricular dysfunction, 7
Combining electrocardiography and necrosis biomarkers improved detection of acute myocardial infarction in patients with LBBB, 29
An early invasive strategy reduced the incidence of major cardiac events in patients with unstable coronary syndromes, 4
Mortality did not vary with sex by 1 year after a first acute myocardial infarction, 31
Reteplase plus abciximab and reteplase alone led to similar 30-day mortality rates in acute MI, 42
Review: Multidisciplinary CHD management programs improve the process of care and reduce hospitalizations, 73
Review: Some new bolus tPA-derived fibrinolytics are as effective as tPA in acute MI, but others increase bleeding, 6
Tenecteplase plus enoxaparin or abciximab was better than tenecteplase plus unfractionated heparin for acute MI, 43
The TIMI risk score predicted mortality in patients with ST-elevation myocardial infarction, 76
- Myocardial ischemia
Clopidogrel plus aspirin was effective but increased bleeding in acute coronary syndromes without ST-segment elevation, 45
Review: Platelet glycoprotein IIb/IIIa blockers for PCI or acute coronary syndromes reduce death and MI but increase bleeding, 41
- Myocardial revascularization
Revascularization led to less angina and fewer adverse cardiac events than did optimal medical care in angina pectoris in the elderly, 47
- Myocardial infarction
Abciximab plus stenting reduced coronary events in patients with acute myocardial infarction, 89
- ### N
- Nadolol
Nadolol and isosorbide prevented recurrent variceal bleeding better than did endoscopic ligation in cirrhosis, 94
- Nausea
Review: Cannabinoids control chemotherapy-induced nausea and vomiting but increase the risk for side effects, 19
- Nebulizers and vaporizers
Review: Pressurized metered-dose inhalers are as effective as other hand-held inhalers for corticosteroid use in asthma, 110
Review: Pressurized metered-dose inhalers are as effective as other hand-held inhalers for β_2 -agonist bronchodilator use in asthma, 111
- Neoplasms
Itraconazole was as effective as amphotericin B for fever and neutropenia in cancer and led to fewer adverse events, 58
Review: Cannabinoids and codeine have similar effects on pain relief, but cannabinoids commonly cause adverse psychotropic effects, 18

Neuraminidase

Oseltamivir was safe and effective for prophylaxis of influenza in the frail elderly, 24

Neutropenia

Itraconazole was as effective as amphotericin B for fever and neutropenia in cancer and led to fewer adverse events, 58

Nurse clinicians

A specialist-nurse intervention reduced readmissions in patients hospitalized with chronic heart failure, 87

O

Obesity

Continuous and intermittent sibutramine were equally effective at 44 weeks for reducing weight in obese persons, 49

Omeprazole

Lansoprazole provided more effective and faster relief for heartburn than did omeprazole in erosive esophagitis, 95

Osteolysis

More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery, 33

Osteoporosis, postmenopausal

Review: Etidronate decreases vertebral but not nonvertebral fractures in postmenopausal women, 98

P

Pain, drug therapy

Review: Cannabinoids and codeine have similar effects on pain relief, but cannabinoids commonly cause adverse psychotropic effects, 18

Parenteral nutrition

Enteral nutrition led to fewer postoperative complications than did parenteral feeding in gastrointestinal cancer, 93

Parkinson disease

Initial bromocriptine did not change mortality in early, mild Parkinson disease, 109

Patient acceptance of health care

Clinical expertise in the era of evidence-based medicine and patient choice (Editorial), A-11 (No. 2)

Patient care management

Review: Multidisciplinary CHD management programs improve the process of care and reduce hospitalizations, 73

Patient satisfaction

Overall satisfaction increased more with inhaled insulin than with subcutaneous insulin in type 1 diabetes, 101

Pectins

A rice-based diet with green banana or pectin reduced diarrhea in infants better than a rice diet alone, 67

Pelvic pain

More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery, 33

Perindopril

Combined therapy with indapamide and perindopril but not perindopril alone reduced the risk for recurrent stroke, 51

Pharyngitis

A clinical prediction model did well in diagnosing pediatric group A β -hemolytic streptococcal pharyngitis, 37

Physical examination

Review: Physical diagnostic tests have low diagnostic accuracy for meniscal lesions of the knee, 113

Physical fitness

Physiotherapy, aerobics, and training devices reduced pain intensity and frequency in chronic low-back pain, 65

Physical therapy

Levodopa plus carbidopa before physiotherapy increased motor recovery after stroke, 52

Physiotherapy, aerobics, and training devices reduced pain intensity and frequency in chronic low-back pain, 65

Placebo effect

Review: Placebo is better than no treatment for subjective continuous outcomes and for treatment of pain, 20

Platelet aggregation inhibitors

Abciximab plus stenting reduced coronary events in patients with acute myocardial infarction, 89

Clopidogrel plus aspirin was effective but increased bleeding in acute coronary syndromes without ST-segment elevation, 45

Platelet glycoprotein GPIIb-IIIa complex

Abciximab did not reduce death or myocardial infarction in the acute coronary syndrome without early revascularization, 44

Abciximab was more effective than tirofiban in preventing ischemic events in patients having coronary stenting, 5

An early invasive strategy reduced the incidence of major cardiac events in patients with unstable coronary syndromes, 4

Reteplase plus abciximab and reteplase alone led to similar 30-day mortality rates in acute MI, 42

Review: Platelet glycoprotein IIb/IIIa blockers for PCI or acute coronary syndromes reduce death and MI but increase bleeding, 41

Tenecteplase plus enoxaparin or abciximab was better than tenecteplase plus unfractionated heparin for acute MI, 43

Point-of-care systems

The TIMI risk score predicted mortality in patients with ST-elevation myocardial infarction, 76

Postoperative complications

Enteral nutrition led to fewer postoperative complications than did parenteral feeding in gastrointestinal cancer, 93

Pregnancy

More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery, 33

Preventive medicine

Rule-based computerized reminders increased ordering of preventive services in an inpatient setting, 74

Primary health care

Review: Primary care counseling improves psychological symptoms in patients with psychological and psychosocial problems, 108

Problem solving

Review: Problem-solving treatment after deliberate self-harm improves depression, hopelessness, and personal problems, 64

Professional competence

Clinical expertise in the era of evidence-based medicine and patient choice (Editorial), A-11 (No. 2)

Prone position

Prone positioning for acute respiratory failure improved short-term oxygenation but not survival, 55

Psychotherapy

Brief psychodynamic interpersonal therapy after deliberate self-poisoning reduced suicidal ideation and deliberate self-harm, 27

Psyllium

Dietary fiber supplementation with psyllium or gum arabic reduced fecal incontinence in community-living adults, 23

Puerperal disorders

More women with the pelvic girdle syndrome than with other pelvic pain during pregnancy had pelvic pain 2 years after delivery, 33

Pulmonary embolism

Helical computed tomographic results indicated small-to-moderate changes in the probability of pulmonary embolism, 30

Pulmonary fibrosis

Clinical and radiologic evaluation had moderate sensitivity and specificity for detecting idiopathic pulmonary fibrosis, 70

R

Radius fractures

Review: Evidence on surgical interventions for distal radial fractures is inconclusive, 112

Receptors, angiotensin

Irbesartan was renoprotective in patients with type 2 diabetes, hypertension, and microalbuminuria, 82

Rehydration solutions

Review: Albumin administration is not associated with excess mortality in acutely ill patients, 2

Reminder systems

Rule-based computerized reminders increased ordering of preventive services in an inpatient setting, 74

Renal artery obstruction

Review: CT angiography and magnetic resonance imaging are the best less-invasive tests for renal artery stenosis, 69

Respiration, artificial

Prone positioning for acute respiratory failure improved short-term oxygenation but not survival, 55

Respiratory distress syndrome, adult

Prone positioning for acute respiratory failure improved short-term oxygenation but not survival, 55

Respiratory tract infections

Zinc or vitamin A reduced diarrhea in young, poor Bangladeshi children, 66

Rice

A rice-based diet with green banana or pectin reduced diarrhea in infants better than a rice diet alone, 67

Risk assessment

The Framingham scores overestimated the risk for coronary heart disease in Japanese, Hispanic, and Native American cohorts, 36

The TIMI risk score predicted mortality in patients with ST-elevation myocardial infarction, 76

Risk factors

The Framingham scores overestimated the risk for coronary heart disease in Japanese, Hispanic, and Native American cohorts, 36

S

Selegiline

Initial bromocriptine did not change mortality in early, mild Parkinson disease, 109

Self-injurious behavior

Brief psychodynamic interpersonal therapy after deliberate self-poisoning reduced suicidal ideation and deliberate self-harm, 27

Review: Problem-solving treatment after deliberate self-harm improves depression, hopelessness, and personal problems, 64

Sepsis

Early goal-directed therapy reduced mortality and multiorgan dysfunction in severe sepsis or septic shock, 90

Sex factors

Mortality did not vary with sex by 1 year after a first acute myocardial infarction, 31

Sigmoidoscopy

Review: Hemoccult screening reduces death from colorectal cancer in average-risk patients > 50 years of age, 91

Socioeconomic factors

The Framingham scores overestimated the risk for coronary heart disease in Japanese, Hispanic, and Native American cohorts, 36

Status epilepticus

Benzodiazepines were as safe as and more effective than placebo for out-of-hospital status epilepticus, 86

Stents

Abciximab plus stenting reduced coronary events in patients with acute myocardial infarction, 89

Streptococcal infections

A clinical prediction model did well in diagnosing pediatric group A β -hemolytic streptococcal pharyngitis, 37

Subclavian vein

Femoral venous catheterization was associated with increased risk for infection and thrombosis in critically ill patients, 1

Suicide

Brief psychodynamic interpersonal therapy after deliberate self-poisoning reduced suicidal ideation and deliberate self-harm, 27

Review: Long-term lithium treatment lowers suicide risk in major affective disorder, 63

Surgery

Review: Evidence on surgical interventions for distal radial fractures is inconclusive, 112

Survival rate

Abciximab did not reduce death or myocardial infarction in the acute coronary syndrome without early revascularization, 44

Syncope

A standardized sequential clinical examination identified probable causes of syncope in 69% of patients, 77

T

Telemedicine

Home monitoring service was more effective than usual care in patients with essential hypertension, 13

Thrombolytic therapy

Review: Thrombolysis increases short-term death and intracranial hemorrhage but decreases long-term death or dependency, 50

Ticlopidine

Clopidogrel plus aspirin was effective but increased bleeding in acute coronary syndromes without ST-segment elevation, 45

Tissue plasminogen activator

Review: Some new bolus tPA-derived fibrinolytics are as effective as tPA in acute MI, but others increase bleeding, 6

TNFR-Fc fusion protein

Review: Adding newer disease-modifying drugs or biological agents to methotrexate improved rheumatoid arthritis symptoms, 97

Tomography, computed

Helical computed tomographic results indicated small-to-moderate changes in the probability of pulmonary embolism, 30

Review: CT angiography and magnetic resonance imaging are the best less-invasive tests for renal artery stenosis, 69

Triamcinolone acetonide

Salmeterol was not as effective as triamcinolone for persistent asthma, 14

Stopping triamcinolone led to treatment failures in patients with persistent asthma who were receiving salmeterol and triamcinolone, 15

V

Vascular diseases

Long-term and short-term oral anticoagulation therapies were equivalent for venous thromboembolism, 10

Venous thrombosis

3 months and 1 year of oral anticoagulant therapy were equivalent for idiopathic proximal deep venous thrombosis, 11

Long-term and short-term oral anticoagulation therapies were equivalent for venous thromboembolism, 10

Review: Extended-duration prophylaxis with heparin prevents deep venous thrombosis in hip or knee replacement, 9

Ventricular dysfunction, left

B-natriuretic peptide levels had high sensitivity and specificity for detecting abnormal ventricular function, 68

Vitamin A

Zinc or vitamin A reduced diarrhea in young, poor Bangladeshi children, 66

Vomiting

Review: Cannabinoids control chemotherapy-induced nausea and vomiting but increase the risk for side effects, 19

W

Waiting lists

A nurse-led intervention reduced risk factors, anxiety, and depression in patients waiting for CABG, 46

Weight loss

Continuous and intermittent sibutramine were equally effective at 44 weeks for reducing weight in obese persons, 49

Z

Zinc

Zinc or vitamin A reduced diarrhea in young, poor Bangladeshi children, 66

